ARUNDEL PEDIATRICS, P.A.

Tylenol (Acetaminophen) Dosage Chart

** Tylenol can be given every 4 HOURS ** DO NOT give to children under 2 MOS OF AGE **

Weight (lbs)	Dose (mgs)	Infant Oral Suspension (160mg/5ml)	Children's Oral Suspension (160mg/5ml)	Chewable (80 mg tab)	Jr. Strength (160 mg tab)	Adult tab 325 mg
6-11 lbs	40 mg	$1.25 \text{ ml} = \frac{1}{4} \text{ tsp}$				
12-17 lbs	80 mg	$2.5 \text{ ml} = \frac{1}{2} \text{ tsp}$	$2.5 \text{ ml} = \frac{1}{2} \text{ tsp}$			
18-22 lbs	120 mg	$3.75 \text{ ml} = \frac{3}{4} \text{ tsp}$	$3.75 \text{ ml} = \frac{3}{4} \text{ tsp}$	1 ½ tabs		
23-35 lbs	160 mg	5 ml = 1 tsp	5 ml =1 tsp	2 tabs	1 tab	
36-47 lbs	240 mg		$7.5 \text{m} = 1\frac{1}{2} \text{tsp}$	3 tabs	1½ tab	
48-59 lbs	320 mg		10 ml = 2 tsp	4 tabs	2 tabs	1 tab
60-71 lbs	400 mg		$12.5 \text{ ml} = 2\frac{1}{2} \text{ tsp}$	5 tabs	2½ tabs	1 tab
72-95 lbs	480 mg		15 ml = 3 tsp	6 tabs	3 tabs	1½ tab
96+ lbs	640 mg		20 ml = 4 tsp	8 tabs	4 tabs	2 tabs

Motrin & Advil (Ibuprofen) Dosage Chart

** Ibuprofen can be given every 6 HOURS ** DO NOT give to children under 6 MOS OF AGE **

	=	-	_			
Weight (lbs)	Dose (mgs)	Infant Oral Suspension (50mg/1.25ml)	Children's Oral Suspension (100mg/5ml)	Chewable (50 mg tab)	Jr. Strength (100 mg tab)	Adult tab 200 mg
6-11 lbs						
12-17 lbs	50 mg	1.25 ml=1 drop	$2.5 \text{ ml} = \frac{1}{2} \text{ tsp}$			
18-22 lbs	75 mg	1.875 ml=1½ drops	$3.75 \text{ ml} = \frac{3}{4} \text{ tsp}$	1 ½ tabs		
23-35 lbs	100 mg	2.5 ml=2 drops	5 ml =1 tsp	2 tabs	1 tab	
36-47 lbs	240 mg		7.5 m l= 1½ tsp	3 tabs	1½ tab	
48-59 lbs	320 mg		10 ml = 2 tsp	4 tabs	2 tabs	1 tab
60-71 lbs	400 mg		$12.5 \text{ ml} = 2\frac{1}{2} \text{ tsp}$	5 tabs	2½ tabs	1 tab
72-95 lbs	480 mg		15 ml = 3 tsp	6 tabs	3 tabs	1½ tab
96+ lbs	640 mg		20 ml = 4 tsp	8 tabs	4 tabs	2 tabs

Benadryl (Diphenhydramine) Dosage Chart

** Benadryl can be given every 6 HOURS ** Consult your physician for children under 12 MOS OF AGE **

Weight (lbs)	Age (years)	Suspension (12.5mg/5ml)	Chewable (12.5 mg tab)	Fastmelts (12.5 mg tab)	Tab/Cap 25 mg
12-19 lbs	1-1½ yrs	$2.5 \text{ ml} = \frac{1}{2} \text{ tsp}$			
20-26 lbs	1½-2 yrs	$3.75 \text{ ml} = \frac{3}{4} \text{ tsp}$			
27-39 lbs	2-4 yrs	5 ml = 1 tsp	1 tab	1 tab	
40-52 lbs	5-6 yrs	$7.5 \text{ ml} = 1 \frac{1}{2} \text{ tsp}$	1 ½ tab	1 ½ tab	
53-67 lbs	7-8 yrs	10 ml = 2 tsp	2 tabs	2 tabs	1 tab/cap
68-79 lbs	9-10 yrs	$10-12.5$ ml = $2-2\frac{1}{2}$ tsp	2-2½ tabs	2 tabs	1 tab/cap
80-95 lbs	11-12 yrs	10-15 ml = 2-3 tsp	2-3 tabs	2-3 tabs	1 tab/cap
96+ lbs	12+ yrs	10-20 ml = 2-4 tsp	2-4 tabs	2-4 tabs	1-2 tabs/caps

^{**} CAUTION!!! Benadryl can cause significant sleepiness or an unexpected hyperactivity reaction in some children **

** Dosing by weight is most accurate ** Consult your physician for any questions **